

PROXEN SR®

(naproxen, sustained release)

NAME OF THE MEDICINE



CAS registry number: 2204531

DESCRIPTION

PROXEN SR (naproxen, sustained release) is a non-steroidal anti-inflammatory drug (NSAID) with analgesic, anti-inflammatory and antipyretic properties.

Naproxen is a propionic acid derivative related to the arylacetic acid class of drugs. It is unrelated to salicylates and the corticosteroid hormones. The chemical name of naproxen is (+)-6-methoxy-alpha-methyl-2-naphthaleneacetic acid. It has a chemical formula of C₁₄H₁₄O₃ and a molecular weight of 230.3. It is an odourless, white to off-white crystalline substance. It is lipid soluble, practically insoluble in water at low pH and freely soluble in water at high pH.

PROXEN SR is available as a sustained release tablet containing 750 mg or 1000 mg of naproxen. PROXEN SR tablets contain the excipients hypromellose, magnesium stearate, Sunset Yellow CI 15985 and purified water.

PHARMACOLOGY

Pharmacodynamics

Naproxen has been shown to have anti-inflammatory properties when tested in human clinical studies. In addition, it has analgesic and antipyretic actions. It exhibits its anti-inflammatory effects even in adrenalectomised animals, indicating that its action is not mediated through the pituitary axis. It inhibits prostaglandin synthetase, as do other NSAIDs, however, the exact mechanism of its anti-inflammatory action is not known.

Pharmacokinetics

Absorption

In humans naproxen is completely absorbed from the gastrointestinal tract after oral administration.

PROXEN SR tablets have a matrix formulation which remains intact during its transit through the gastrointestinal tract. When the tablet comes in contact with gastrointestinal fluids it forms a gelatinous layer which controls the rate of release of naproxen. Due to the controlled release, the peak plasma levels of naproxen are simultaneously reduced and delayed compared to standard NAPROSYN tablets, although the mean area under the plasma concentration-time curve of the SR formulation is equivalent to that of the standard formulation.

The lowest plasma naproxen concentrations attained with PROXEN SR tablets given once daily (i.e. 750 mg once daily or 1000 mg once daily) are equivalent to the lowest plasma naproxen concentration attained with the corresponding standard NAPROSYN tablet formulations given twice daily (i.e. 375 mg twice daily or 500 mg twice daily).

PROXEN SR tablets are unique compared to the standard NAPROSYN tablet in that the 'peak' to 'trough' plasma naproxen concentration ratio of 2.6 for NAPROSYN is reduced to 1.8 for the sustained release formulation, hence indicating that the administration of PROXEN SR results in a more favourable pharmacokinetic profile.

Food does not alter the bioavailability of PROXEN SR tablets. However, ingestion of a high-fat meal prior to administration of PROXEN SR tablets does significantly increase the peak plasma concentration compared to that seen in fasted subjects. Food decreases the rate of transit of PROXEN SR tablets from the stomach to the intestines in both young and old adult patients. These results are unlikely to be of clinical significance.

Distribution

Naproxen has a relatively small volume of distribution (0.09 + 0.03 L/kg), which corresponds to about 10% of the body weight in humans. At therapeutic levels naproxen is greater than 99% albumin-bound.

The plasma concentration of naproxen increases proportionally with doses up to 500 mg twice daily. Larger doses result in a less than proportional increase due to accelerated renal clearance of disproportionately increased amounts of non-protein bound drug. However, whether this effect increases or decreases the toxicity of naproxen has not been established.

Steady-state plasma levels of naproxen are reached after 4 to 5 doses.

Naproxen enters synovial fluid and crosses the placenta. It has been found in the milk of lactating mothers at a concentration approximately 1% of that found in plasma.

Metabolism

Naproxen is metabolised in the liver to 6-O-desmethyl naproxen (approximately 28% of an IV dose).

Elimination

Approximately 95% of the naproxen is excreted in the urine, primarily as naproxen (10%), 6-O-desmethyl naproxen (5%) or their conjugates. The rate of excretion of metabolites and conjugates has been found to coincide closely with the rate of naproxen clearance from the plasma. Small amounts, 5% or less, are excreted in the faeces.

The elimination half-life of naproxen is approximately 14 hours.

Pharmacokinetics in Special Populations

Children

The pharmacokinetic profile of naproxen in children aged 5-16 years is similar to that in adults.

Renal Impairment

Given that naproxen and its metabolites are primarily excreted by the kidney, the potential exists for accumulation in the presence of renal insufficiency. Elimination of naproxen is decreased in patients with severe renal impairment (creatinine clearance < 20 mL/min), in whom there is higher clearance of naproxen than estimated from the degree of renal impairment alone (see PRECAUTIONS – Renal Impairment).

INDICATIONS

PROXEN SR is indicated for the treatment of rheumatoid arthritis, osteoarthritis, ankylosing spondylitis and for the relief of chronic pain states in which there is an inflammatory component.

CONTRAINDICATIONS

PROXEN SR is contraindicated in patients:

- who are hypersensitive to naproxen or naproxen sodium or in whom acetylsalicylic acid (aspirin) or other non-steroidal anti-inflammatory/analgesic agents induce allergic manifestations, e.g. asthma, nasal polyps, rhinitis and urticaria. Severe anaphylactic-like reactions to naproxen have been reported in such patients
- with either active, or a history of, peptic or gastrointestinal ulceration, chronic dyspepsia or active gastrointestinal bleeding or perforation, related to previous NSAIDs therapy
- with active, or history of recurrent peptic ulcer/haemorrhage (two or more distinct episodes of proven ulceration or bleeding)
- in patients less than 2 years of age since safety in this age group has not been established
- with severe heart failure
- undergoing treatment of perioperative pain in setting of coronary artery surgery (CABG)
- with severe hepatic impairment

PRECAUTIONS

Cardiovascular Thrombotic Events

Observational studies have indicated that non-selective NSAIDs may be associated with an increased risk of serious cardiovascular events including myocardial infarction and stroke, which may increase with dose or duration of use. Patients with cardiovascular disease, history of atherosclerotic cardiovascular disease or cardiovascular risk factors may also be at greater risk. To minimise the potential risk of an adverse cardiovascular event in patients taking an NSAID, especially in those with cardiovascular risk factors, the lowest effective dose should be used for the shortest possible duration (see DOSAGE AND ADMINISTRATION).

Physicians and patients should remain alert for such CV events even in the absence of previous CV symptoms. Patients should be informed about signs and/or symptoms of serious CV toxicity and the steps to take if they occur.

There is no consistent evidence to suggest that concurrent use of aspirin mitigates the possible increased risk of serious cardiovascular thrombotic events associated with NSAID use.

Clinical trial and epidemiological data suggest that use of coxibs and some NSAIDs (particularly at high doses and long term treatment) may be associated with a small increased risk of arterial thrombotic events (e.g. myocardial infarction or stroke).

Hypertension

NSAIDs may lead to onset of new hypertension or worsening of pre-existing hypertension and patients taking anti-hypertensives with NSAIDs may have an impaired anti-hypertensive response. Caution is advised when prescribing NSAIDs to patients with hypertension. Blood pressure should be monitored closely during initiation of NSAID treatment and at regular intervals thereafter.

Heart Failure

Fluid retention and oedema have been observed in some patients taking NSAIDs, therefore caution is advised in patients with fluid retention or heart failure.

Gastrointestinal

All NSAIDs can cause gastrointestinal discomfort and rarely serious, potentially fatal, gastrointestinal effects such as ulcers, irritation, bleeding and perforation which may increase with dose or duration of use, but can occur at any time without warning. Upper gastrointestinal ulcers, gross bleeding or perforation caused by NSAIDs occur in approximately 1% of patients treated for 3 - 6 months and in about 2 - 4% of patients treated for one year. These trends continue with longer duration of use, increasing the likelihood of developing a serious gastrointestinal event at some time during the course of therapy. However, even short term therapy is not without risk.

Caution is advised in patients with risk factors for gastrointestinal events who may be at greater risk of developing serious gastrointestinal events e.g. elderly, debilitated patients, those with a history of serious gastrointestinal events, smoking and alcoholism.

NSAIDs should be given with care to patients with a history of inflammatory bowel disease (ulcerative colitis; Crohn's disease) as their condition may be exacerbated. Patients with a history of gastrointestinal toxicity, particularly when elderly, should report any unusual abdominal symptoms (especially gastrointestinal bleeding) particularly in the initial stages of treatment. When gastrointestinal bleeding or ulceration occurs in patients receiving NSAIDs, treatment should be withdrawn immediately. Physicians should warn patients about the signs and symptoms of serious gastrointestinal toxicity.

Studies to date have not identified any subset of patients not at risk of developing peptic ulcer and bleeding. However, the elderly have an increased frequency of adverse effects to NSAIDs, especially gastrointestinal bleeding and perforation which may be fatal. Debilitated patients do not seem to tolerate ulceration or bleeding as well as others. Most of the fatal gastrointestinal events associated with NSAIDs occurred with the elderly and/or debilitated patients.

In patients with active peptic ulcer or inflammatory disease of the gastrointestinal tract and rheumatoid arthritis, an attempt might be made to treat the arthritis with a non-ulcerogenic drug.

Caution is advised in patients receiving concomitant medications which could increase the risk of ulceration or bleeding (see PRECAUTIONS - Interactions with Other Medicines). The concurrent use of aspirin and NSAIDs also increase the risk of serious gastrointestinal adverse events.

Patients with risk factors should commence treatment on the lowest dose available.

Use in Renal Impairment

There have been reports of impaired renal function, renal failure, acute interstitial nephritis, haematuria, proteinuria, renal papillary necrosis, and occasionally nephritic syndrome associated with PROXEN SR.

PROXEN SR should not be given to patients with creatinine clearance less than 30 mL/minute because accumulation of naproxen metabolites has been seen in such patients.

As with other NSAIDs, PROXEN SR should be used with caution in patients with impaired renal function, or a history of kidney disease because naproxen is an inhibitor of prostaglandin synthesis. Caution should be observed in patients with conditions leading to a reduction in blood volume and/or renal blood flow as prostaglandins have a supportive role in the maintenance of renal perfusion. In these patients administration of PROXEN SR or other NSAIDs may cause a dose-dependant reduction in renal prostaglandin formation and may precipitate overt renal decompensation or failure. Patients at greatest risk are those with impaired renal function, hypovolaemia, heart failure, liver dysfunction, salt depletion, those taking diuretics, angiotensin converting enzyme inhibitors or angiotensin receptor blockers and the elderly. Discontinuation of PROXEN SR is usually followed by recovery to the pre-treatment state; however, serious adverse events may persist. PROXEN SR should be used with great caution in such patients and the monitoring of serum creatinine and/or creatinine clearance is advised and patients should be adequately hydrated. A reduction of daily dosage should be considered to avoid the possibility of excessive accumulation of naproxen metabolites in these patients.

Haemodialysis does not decrease the plasma concentration of naproxen because of the high degree of its protein binding.

Haematological

Naproxen decreases platelet aggregation and prolongs bleeding time. This effect should be kept in mind when bleeding times are being determined (see PRECAUTIONS – Effects on Laboratory Tests).

Patients who have coagulation disorders or are receiving drug therapy that interferes with haemostasis should be carefully observed if PROXEN SR is administered. Patients at high risk of bleeding and those on anticoagulation therapy (e.g. heparin or dicoumarol derivatives) may be at increased risk of bleeding if given PROXEN SR concurrently. Therefore, the benefits of prescribing PROXEN SR should be weighed against these risks.

Patients with initial haemoglobin values of 10 grams or less, and who are to receive long-term therapy should have haemoglobin values determined frequently.

Patients on other drugs such as hydantoins, sulfonamides, sulfonyleureas or methotrexate should be observed for increased effect or toxicity (see PRECAUTIONS – Interactions with Other Medicines).

Severe Skin Reactions

NSAIDs may very rarely cause serious cutaneous adverse events such as exfoliative dermatitis, Stevens-Johnson Syndrome (SJS) and toxic epidermal necrolysis (TEN), which can be fatal and occur without warning. These serious adverse events are idiosyncratic and are independent of dose or duration of use. Patients should be advised of the signs and symptoms of serious skin reactions and to consult their physician at the first appearance of a skin rash or any other sign of hypersensitivity.

Anaphylactic Reactions

Hypersensitivity reactions may occur in susceptible individuals. Anaphylactic (anaphylactoid) reactions may occur both in patients with, and without, a history of hypersensitivity or exposure to aspirin or other NSAIDs or naproxen-containing products. They may also occur in individuals with a history of angioedema, bronchospastic reactivity (e.g. asthma), rhinitis and nasal polyps. Anaphylactoid reactions, like anaphylaxis, may have a fatal outcome.

Bronchospasm may be precipitated in patients suffering from, or with a history of, asthma or allergic disease or aspirin sensitivity.

Use in Hepatic Impairment

As with other NSAIDs borderline elevations of one or more liver function tests may occur in up to 15% of patients. These abnormalities may progress, may remain essentially unchanged, or may resolve with continued therapy. The ALT test is probably the most sensitive indicator of liver dysfunction. Meaningful elevations (three times the upper limit of normal) of ALT or AST occurred in controlled clinical trials in less than 1% of patients. Physicians and patients should remain alert for hepatotoxicity. Patients should be

informed about the signs and/or symptoms of hepatotoxicity. A patient with symptoms and/or signs suggesting hepatic dysfunction (e.g. nausea, fatigue, lethargy, pruritis, jaundice, abdominal tenderness in the right upper quadrant and “flu-like” symptoms), or in whom an abnormal hepatic test has occurred, should be evaluated for evidence of the development of more severe hepatic reaction while on therapy with PROXEN SR.

Hepatic abnormalities may be the result of hypersensitivity or direct toxicity.

Severe hepatic reactions, including jaundice and cases of fatal hepatitis, have been reported with naproxen as with other NSAIDs. Cross reactivity has been reported. Although such reactions are rare, if abnormal hepatic tests persist or worsen, if clinical signs and symptoms consistent with hepatic disease develop, or if systemic manifestations occur (e.g. eosinophilia, rash, etc.), PROXEN SR should be discontinued.

Chronic alcoholic hepatic disease and potentially other forms of cirrhosis reduce the total plasma concentration of naproxen; however, the plasma concentration of unbound naproxen is increased. The implication of this finding for naproxen dosing is unknown.

In patients with impaired hepatic function, the lowest effective dose is recommended.

Infection

The antipyretic, anti-inflammatory and analgesic effects of naproxen may mask the usual signs or symptoms of infection.

Ocular Events

Adverse ophthalmological effects have been observed with NSAIDs. In rare cases, adverse ocular disorders including papillitis, retrobulbar optic neuritis and papilloedema have been reported in users of NSAIDs including PROXEN SR, although a cause-and-effect relationship cannot be established; accordingly, patients who develop visual disturbances during treatment with PROXEN SR should have an ophthalmological examination.

Fluid Retention and Oedema

Peripheral oedema has been observed in some patients taking PROXEN SR or other NSAIDs. Although sodium retention has not been reported in metabolic studies, it is possible that patients with compromised cardiac function may be at greater risk when taking naproxen. For this reason, naproxen should be used with caution in patients with fluid retention, hypertension or heart failure.

Use in Pregnancy

PREGNANCY CATEGORY: C

NSAIDs inhibit prostaglandin synthesis and when given during the latter part of pregnancy, may cause closure of the foetal ductus arteriosus, prolong labour and delay birth. During the last few days before expected birth, agents with an inhibitory effect on prostaglandin synthesis should be avoided. Continuous treatment with NSAIDs during the last month of pregnancy should only be given when clearly indicated.

PROXEN SR should only be administered during pregnancy if the benefit justifies the potential risk.

The use of PROXEN SR, as with any drug known to inhibit cyclooxygenase/prostaglandin synthesis, may impair fertility and is not recommended in women attempting to conceive. In women who have difficulty conceiving or are undergoing investigation of infertility, withdrawal of naproxen should be considered.

Data from epidemiological studies suggest an increased risk of miscarriage after the use of a prostaglandin synthesis inhibitor in early pregnancy.

Use in Lactation

Naproxen has been found in the milk of lactating mothers at a concentration approximately 1% of that found in plasma. As the effect of naproxen in the newborn is not known, the use of PROXEN SR in lactating mothers is not recommended.

Paediatric Use

PROXEN SR is not recommended in children under 5 years of age as the safety and efficacy in this population has not been established.

Use in Elderly

The lowest effective dose is recommended in elderly patients.

Studies indicate that although the total plasma concentration of naproxen is unchanged, the unbound plasma fraction of naproxen is increased in the elderly.

PROXEN SR is not recommended in elderly patients with hepatic or renal impairment.

INTERACTIONS WITH OTHER MEDICINES

Concomitant administration of sucralfate or cholestyramine can delay the absorption of naproxen, but does not affect its extent. Antacids have a variable effect on absorption.

Other NSAIDs

Combination of naproxen-containing products and other NSAIDs, including cyclooxygenase-2 (COX-2) selective inhibitors, is not recommended, because of the cumulative risks of inducing serious NSAID-related adverse events.

Protein Binding

Naproxen is highly bound to plasma albumin; thus naproxen has a theoretical potential for interaction with other albumin-bound drugs, for example, warfarin or bishydroxycoumarin may be displaced and induce excessively prolonged prothrombin times. Similarly, patients receiving hydantoin, sulfonamides or sulfonylureas should be observed for increased effect or toxicity (see PRECAUTIONS – Haematological).

Warfarin

The concurrent use of NSAIDs and warfarin has been associated with severe, sometimes fatal, haemorrhage. The exact mechanism of the interaction between NSAIDs and warfarin is unknown, but may involve enhanced bleeding from NSAID-induced gastrointestinal ulceration, or an additive effect of anticoagulation by warfarin and inhibition of platelet function by NSAIDs. PROXEN SR should be used in combination with warfarin only if absolutely necessary, and patients taking this combination of drugs should be closely monitored.

Anticoagulant/ Anti-platelet Agents

Patients who have coagulation disorders or are receiving drug therapy that interferes with haemostasis should be carefully observed if naproxen is administered. Patients on full anticoagulation therapy (e.g. heparin or dicoumarol derivatives) may be at increased risk of bleeding if given naproxen concurrently. Thus, the benefits should be weighed against these risks.

There is an increased risk of gastrointestinal bleeding when anti-platelet agents are combined with NSAIDs.

Selective Serotonin Reuptake Inhibitors (SSRIs)

There is an increased risk of gastrointestinal bleeding when SSRIs are combined with NSAIDs.

Steroids

If steroid dosage is reduced or eliminated during PROXEN SR therapy, the steroid dosage should be reduced slowly and the patient must be observed closely for any evidence of adverse effects, including adrenal insufficiency and exacerbation of symptoms of underlying disease.

Probenecid

Probenecid significantly prolongs the half-life of naproxen (from 14 to 37 hrs). This is associated with a decrease in conjugated metabolites and an increase in 6-O-desmethyl naproxen.

Methotrexate

Concomitant administration of naproxen and methotrexate should be administered with caution, because naproxen has been reported among other NSAIDs to reduce the tubular secretion of methotrexate in animal models, and thus possibly increasing the toxicity of methotrexate.

Beta-Blockers

Naproxen and other NSAIDs can reduce the anti-hypertensive effect of beta-blockers, angiotensin converting enzyme inhibitors (ACE inhibitors), and angiotensin receptor blockers (ARBs).

Diuretics

As with other NSAIDs, naproxen may inhibit the natriuretic effect of furosemide.

Lithium

Inhibition of renal lithium clearance leading to increases in plasma lithium concentrations has been reported.

Sodium Bicarbonate

Sodium bicarbonate may enhance the rate of naproxen absorption.

Zidovudine

In vitro studies have shown that naproxen may interfere with the metabolism of zidovudine, resulting in higher zidovudine plasma levels. Therefore, to avoid the potential side effects associated with increased zidovudine plasma levels, dose reduction should be considered.

ACE-Inhibitors

Concomitant use of NSAIDs with ACE inhibitors or angiotensin receptor blockers may increase the risk of renal dysfunction, especially in patients with pre-existing poor renal function (see PRECAUTIONS).

Combination use of ACE-inhibitors or angiotensin receptor antagonists, anti-inflammatory drugs and thiazide diuretics

The use of an ACE inhibiting drug (ACE-inhibitor or angiotensin receptor antagonist), an anti-inflammatory drug (NSAID or COX-2 inhibitor) and a thiazide diuretic at the same time (triple whammy) increases the risk of renal impairment. This includes use in fixed-combination products containing more than one class of drug. Combined use of these medications should be accompanied by increased monitoring of serum creatinine, particularly at the initiation of the combination. The combination of drugs from these three classes should be used with caution particularly in elderly patients or those with pre-existing renal impairment.

Effects on Laboratory Tests

Naproxen decreases platelet aggregation and prolongs bleeding time. This effect should be considered when bleeding times are determined.

PROXEN SR may artefactually interfere with some tests for 17-ketogenic steroid and may interfere with some urinary assays for 5-hydroxy-indoleacetic acid (5HIAA). 17-hydroxycorticosteroid measurements (Porter/Silber test) do not appear to be altered.

Naproxen therapy should be temporarily discontinued for at least 72 hours before testing adrenal function.

Effects on Ability to Drive and Operate Machinery

Some patients may experience drowsiness, dizziness, vertigo, insomnia or depression with the use of PROXEN SR. If patients experience these or similar undesirable effects, they should exercise caution in carrying out activities that require alertness.

ADVERSE EFFECTS

Adverse effects reported in controlled clinical trials in 960 patients treated for rheumatoid arthritis and osteoarthritis are listed below. In general, these effects were reported 2 to 10 times more frequently than they were in studies of 962 patients treated for mild to moderate pain.

Incidence between 3% and 9%

Gastrointestinal: The most frequently reported adverse effects were related to the gastrointestinal tract. These were: constipation, heartburn, abdominal pain, nausea.

Central Nervous System: headache, dizziness, drowsiness

Dermatologic: itching (pruritis), skin eruption, ecchymoses

Special Senses: tinnitus

Cardiovascular: oedema, dyspnoea

Incidence between 1% and less than 3%

Gastrointestinal: dyspepsia, diarrhoea, stomatitis

Central Nervous System: light-headedness, vertigo

Dermatologic: sweating, purpura

Special Senses: hearing disturbances, visual disturbances

Cardiovascular: palpitations

General: thirst

Incidence less than 1%

PROBABLE CAUSAL RELATIONSHIP:

The following adverse effects were reported less frequently than 1% during controlled clinical trials and through post-marketing reports. A causal relationship probably exists between naproxen and these adverse effects.

Gastrointestinal: abnormal liver function tests, gastrointestinal bleeding, haematemesis, jaundice, melaena, peptic ulceration with bleeding and/or perforation, non-peptic gastrointestinal ulceration, vomiting, ulcerative stomatitis, colitis, fatal hepatitis

Renal: glomerular nephritis, haematuria, interstitial nephritis, nephrotic syndrome, renal disease, renal papillary necrosis, renal failure, hyperkalaemia

Haematologic: eosinophilia, granulocytopenia, leukopenia, thrombocytopenia

Central Nervous System: depression, dream abnormalities, inability to concentrate, insomnia, malaise, myalgia, muscle weakness, aseptic meningitis

Dermatologic: porphyria cutanea tarda, epidermolysis bullosa, alopecia, skin rashes, epidermal necrolysis, erythema multiforme, Stevens-Johnson syndrome (SJS), photosensitivity reactions including rare cases in which the skin resembles porphyria cutanea tarda (pseudoporphyria) or epidermolysis bullosa

Special Senses: hearing impairment

Cardiovascular: vasculitis, congestive heart failure

General: menstrual disorders, pyrexia (chills and fever), eosinophilic pneumonitis, anaphylactoid reactions (see PRECAUTIONS – Anaphylactic Reactions)

CAUSAL RELATIONSHIP UNKNOWN:

Other reactions have been reported in circumstances in which a causal relationship could not be established. Although rarely reported, the physician should be alerted to these.

Haematologic: agranulocytosis, aplastic anaemia, haemolytic anaemia

Central and Peripheral Nervous System: cognitive dysfunction, convulsions, paraesthesia

Dermatologic: urticaria, photosensitivity

Mouth and Throat: sore throat

General: angioneurotic oedema, hyperglycaemia, hypoglycaemia, hyperkalaemia

Reproductive: female infertility

Post-Marketing Experience

The following adverse effects have been reported with NSAIDs and PROXEN SR:

Gastrointestinal: inflammation, peptic ulcers, ulceration, perforation and obstruction of the upper and lower gastrointestinal tract, gastrointestinal bleeding (sometimes fatal, particularly in the elderly), heartburn, nausea, Oesophagitis, vomiting, diarrhoea, flatulence, constipation, dyspepsia, abdominal pain, non-peptic gastrointestinal ulceration, melaena, haematemesis, stomatitis, ulcerative stomatitis, exacerbation of ulcerative colitis and Crohn's disease, pancreatitis, gastritis

Infection: aseptic meningitis

Blood and Lymphatic System Disorders: agranulocytosis, aplastic anaemia, eosinophilia, haemolytic anaemia, leucopenia, thrombocytopenia

Immune System Disorders: anaphylactoid reactions

Metabolic and Nutrition Disorders: hyperkalaemia

Psychiatric Disorders: depression, dream abnormalities, insomnia

Nervous System Disorders: dizziness, drowsiness, headache, light-headedness, retrobulbar optic neuritis, convulsions, cognitive dysfunction, inability to concentrate

Eye Disorders: visual disturbances, corneal opacity, papillitis, papilloedema

Ear and Labyrinth Disorders: hearing impairment, hearing disturbances, tinnitus, vertigo

Cardiac Disorders: palpitations, cardiac failure, congestive heart failure

Vascular Disorders: hypertension, vasculitis

Respiratory, Thoracic and Mediastinal Disorders: dyspnoea, pulmonary oedema, asthma, eosinophilic pneumonitis

Hepatobiliary Disorders: hepatitis, jaundice

Skin and Subcutaneous Tissue Disorder: ecchymoses, itching (pruritus), purpura, skin eruptions, sweating, alopecia, epidermal necrolysis, very rarely toxic epidermal necrolysis (TEN), erythema multiforme, bullous reactions (including SJS), erythema nodosum, fixed drug eruption, lichen planus, pustular reaction, skin rashes, systemic lupus erythematosus (SLE), urticaria, photosensitivity reactions, including rare cases resembling porphyria cutanea tarda (pseudoporphyria) or epidermolysis bullosa, or angioneurotic oedema

If skin fragility, blistering or other symptoms suggestive of pseudoporphyria occur, treatment should be discontinued and patient monitored.

Musculoskeletal and Connective Tissue Disorders: myalgia, muscle weakness

Renal and Urinary Disorders: haematuria, interstitial nephritis, nephritic syndrome, renal disease, renal failure, renal papillary necrosis

Reproductive System: female infertility

General Disorders: oedema, thirst

Investigations: abnormal liver function tests, raised serum creatinine

DOSAGE AND ADMINISTRATION

After assessing the risk/benefit ratio in each individual patient, the lowest effective dose for the shortest possible duration should be used.

The total required daily dose of naproxen can be achieved by administering the appropriate single daily dose i.e. PROXEN SR 750 mg or PROXEN SR 1000 mg.

The total daily dose of naproxen should not exceed 1000 mg. PROXEN SR tablets should be taken whole and not chewed.

Patients on long term treatment should be reviewed regularly with regards to efficacy, risk factors and ongoing need for treatment.

PROXEN SR TABLETS ARE NOT INTENDED FOR PATIENTS REQUIRING SHORT-TERM TREATMENT FOR ACUTE INDICATIONS.

OVERDOSAGE

Significant overdose of the medicine may be characterised by dizziness, drowsiness, epigastric pain, abdominal discomfort, indigestion, transient alterations in liver function, hypoprothrombinaemia, renal dysfunction, metabolic acidosis, apnoea, disorientation, nausea or vomiting. A few patients have experienced seizures, but it is unclear if these were causally related to naproxen. It is not known what dose of naproxen would be life-threatening.

Gastrointestinal bleeding may occur. Hypertension, acute renal failure, respiratory depression and coma may occur after the ingestion of NSAIDs, and may occur following an overdose.

Anaphylactoid reactions have been reported with therapeutic ingestion of NSAIDs, and may occur following an overdose.

Patients should be managed by symptomatic and supportive care following NSAIDs overdose. There are no specific antidotes. Prevention of further absorption (e.g. activated charcoal) may be indicated in symptomatic patients seen within 4 hours of ingestion or following a large overdose. Forced diuresis, alkalization of urine, haemodialysis, or haemoperfusion may not be useful due to high protein binding.

Due to the sustained-release characteristic of PROXEN SR tablets, it should be expected that naproxen will continue to be absorbed for up to 16 hours after ingestion.

Contact the Poisons Information Centre for advise on management of overdosage.

PRESENTATION AND STORAGE CONDITIONS

PROXEN SR 750 mg sustained release tablets are available as uncoated, oval, peach coloured tablets, embossed "NPR SR-750" on one side, in PVC/aluminium blister packs of 28s.

PROXEN SR 1000 mg sustained release tablets are available as uncoated, oval, peach coloured tablets, embossed "NPR SR-1000" on one side, in PVC/aluminium blister packs of 28s.

Store below 30°C. Protect from light.

NAME AND ADDRESS OF THE SPONSOR

atnahs Pharma Australia Pty Ltd

Level 10 / 10 Shelley Street,

SYDNEY,

NSW, 2000, Australia

POISON SCHEDULE OF THE MEDICINE

Schedule 4 – Prescription Only Medicine

DATE OF FIRST INCLUSION IN THE AUSTRALIAN REGISTER OF THERAPEUTIC GOODS (THE ARTG):

7 February 1994

DATE OF MOST RECENT AMENDMENT:

23 May 2017